

The brightest stars of the σ Orionis cluster

J. A. Caballero^{1,2,3,*}

¹ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

² Isaac Newton Group, Apartado 321, E-38700 Santa Cruz de La Palma, Canary Islands, Spain

³ Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

Received 27 October 2006; accepted 2 January 2007

ABSTRACT

Context. The very young σ Orionis cluster (~ 3 Ma) is a cornerstone for the understanding of the formation of stars and substellar objects down to planetary masses. However, its stellar population is far to be completely known.

Aims. This study has the purpose of identifying and characterising the most massive stars of σ Orionis to complement current and future deep searches for brown dwarfs and planetary-mass objects in the cluster.

Methods. I have cross-correlated the sources in the Tycho and 2MASS catalogues in a region of 30 arcmin radius with centre in the O-type star σ Ori A. In the area, I have studied the membership in the Ori OB 1b Association of the brightest stars in the optical using astrometric, X-ray and infrared and optical photometric data from public catalogues and spectroscopic data from the literature.

Results. A list of 26 young stars, four candidate young stars and 16 probable foreground stars has arisen from the study. Seven young stars probably harbour discs (four are new). There is no mass-dependence of the disc frequency in the cluster. I have derived for the first time the mass spectrum in σ Orionis from 1.1 to $24 M_{\odot}$ ($\alpha = +2.0^{+0.2}_{-0.1}$; roughly Salpeter-like). I have also provided additional proofs on the existence of several spatially superimposed stellar populations in the direction of σ Orionis. Finally, the cluster may be closer and older than previously considered**.

Key words. open clusters and associations: individual: σ Orionis — stars: general — planetary systems: protoplanetary discs — astronomical data bases: miscellaneous

1. Introduction

The constellation of Orion, the Hunter, is dominated by the stellar components of the Ori OB 1 complex. The complex is probably related to the Barnard Loop, a huge feature with H α in emission that extends up to Eridanus, and to a hole in the distribution of gas H I, which may have been originated from several supernova events that happened about 4 Ma ago (Brown, de Geus & de Zeeuw 1994). The very young Orion complex is a set of four subgroups or associations of very early type stars, named Ass Ori OB 1a, 1b, 1c and 1d. Each of them possesses slightly different locations, ages (1–10 Ma) and heliocentric distances (300–500 pc). Blaauw (1964), Warren & Hesser (1978), Goudis (1982), de Geus et al. (1990), Brown et al. (1994) and other authors have described the boundaries between subgroups and their characteristics.

The σ Orionis cluster lies within the Ori OB 1b Association, also known as the Orion Belt. This association contains three famous perfectly aligned very bright stars: Alnitak (ζ Ori), Alnilam (ϵ Ori) and Mintaka (δ Ori). It has been suggested that the age, distance and radial velocity of the stellar components vary across the association (Hardie et al. 1964; Warren & Hesser 1977a,b; Guetter 1981; Giesekeing 1983; Genzel & Stutzki 1989). The east part of the Ori OB 1b Association, with Alnitak, the Horsehead Nebula, the Flame Nebula (the Orion B cloud, associated to NGC 2024) and the H II region IC 434, would be the farthest and youngest one. Actually, the star formation is still underway close to the Horsehead Nebula, which

mane is illuminated by the σ Ori star¹, the fourth brightest star in the Orion Belt.

Garrison (1967) and Lyngå (1981) were the first investigators to recognize the σ Orionis cluster. It went unnoticed until Wolk (1996) and Walter, Wolk & Sherry (1998) detected and over-density of X-ray sources and a population of pre-main-sequence low-mass stars surrounding the centre of the cluster (see also: Walter et al. 1997; Wolk & Walter 2000). Although it is not as extremely young as the Trapezium cluster in the Ori OB 1d Association (the Orion Sword) and is not as nearby to the Sun as the rich star-forming regions in the Southern Hemisphere (i.e. Chamaeleon, Ophiuchus), the σ Orionis cluster has the advantage of possessing simultaneously a moderate youth, a relative short heliocentric distance and, especially, a very low interstellar extinction (Lee et al. 1968; Béjar et al. 2004b). This fact may be due to strong winds, ultraviolet radiation and/or turbulence generated by the O-type star in the cluster centre, which would sweep away the gas and dust. Furthermore, several authors have suggested that σ Orionis is in reality an H II region (e.g. Reipurth et al. 1998).

There are different determinations of the age of the σ Orionis cluster, based on different methodologies: (i) from the comparison between mid-resolution real and theoretical spectra surrounding the Li I $\lambda 6707.8$ Å youth-indicator line, (ii) from the presence of very early-type stars, (iii) from the disc frequency around stars and (iv) from fits to theoretical isochrones in colour-magnitude diagrams. All the determinations are directed towards a narrow age interval, between 0.5 and 8 Ma. In Table 1, I sum-

* Alexander von Humboldt Fellow at the Max-Planck-Institut für Astronomie, e-mail: caballero@mpia.de.

¹ Through this paper and for clarity, I am using the full and shortened names for stars and clusters, respectively; i.e. the σ Ori star system, the σ Orionis cluster.

Table 1. Different determinations of the age of the σ Orionis cluster.

Age (Ma)	Reference
8	Blaauw (1964)
$<5.1^a$	Warren & Hesser (1978)
7	Blaauw (1991)
1.7 ± 1.1	Brown et al. (1994)
$4.2^{+2.7}_{-1.5}$	Oliveira et al. (2002)
$2-4^b$	Zapatero Osorio et al. (2002b)
2.5 ± 0.3	Sherry, Walter & Wolk (2004)
3.5 ± 3.0	Hernández et al. (2005)
4–6	Briceño et al. (2005)
6.6	Kharchenko et al. (2005)
$\leq 4-6^c$	Sacco et al. (2006)

^a Age not rotation-corrected (upper limit: 5.1 Ma).

^b Upper limit: 8 Ma; lower limit: 1 Ma.

^c They also found three low-mass high probability members that are older than ~ 10 Ma.

marize several age determinations found in the literature. The most probable age of the cluster is about 3 Ma, with conservative -2 and $+5$ Ma error-bars.

The heliocentric distance to σ Ori from the trigonometric parallax measured by the Hipparcos mission is 352^{+166}_{-168} pc. There are also different determinations of the heliocentric distance to the Ori OB 1b Association and to several σ Orionis members (Brown et al. 1994, 1999; Perryman et al. 1997; de Zeeuw et al. 1999; Brown et al. en prep.). Published distances vary between 360^{+70}_{-60} pc and 473 ± 33 pc. The most recent determinations of the heliocentric distance to the σ Orionis cluster were shown by Hernández et al. (2005), who determined the distances from parallax, 443 ± 16 pc, and from colour-magnitude diagrams, 392 ± 20 pc, and by Sherry, Walter & Wolk (2004), who determined an heliocentric distance of 440 ± 40 pc. I will use the value measured by Brown et al. (1994) of 360^{+70}_{-60} pc.

Béjar et al. (1999) opened the era the deep photometric searches in the cluster and detected the first substellar objects in σ Orionis (they have not stable hydrogen burning in their cores, in contrast to stars). Posterior studies have gone deeper, discovering a wealthy population of brown dwarfs (Béjar et al. 2001, 2004b; Scholz & Eislöffel 2004; Caballero et al. 2004; Kenyon et al. 2005) and even objects with masses in the planetary domain (below the deuterium burning mass limit) (Zapatero Osorio et al. 2000; González-García et al. 2006; Caballero et al. 2007a). The σ Orionis cluster is to date the star-forming region with the largest amount of candidate planetary-mass objects (27). Out of them, 12 have low-resolution spectroscopy (Martín et al. 2001; Barrado y Navascués et al. 2001) and three harbour surrounding discs (Caballero et al. 2007b). The cluster has now turned out to be not only the region with the largest stellar density in the Ori OB 1b Association, but one of the most interesting regions in the Milky Way to study and understand the formation of stars, brown dwarfs and, especially, isolated planetary-mass objects.

In spite of the importance of the cluster, its stellar population is paradoxically poorly known. General studies in the Ori OB 1b Association (e.g. Warren & Hesser 1977a, 1977b, 1978), very wide searches with prism-objective and Schmidt plates (Haro & Moreno 1953; Kogure et al. 1989; Wiramihardja et al. 1989, 1991; Nakano et al. 1995) and the ROSAT-based search by Wolk (1996) provided most of the known cluster members at the end

of last century. However, these surveys were obviously biased towards the detection of active Ori OB1 1b stellar members. Some efforts have been recently carried out to improve our knowledge of the stellar population in σ Orionis. Nonetheless, the bias towards H α emitters (Weaver & Babcock 2004), large-amplitude photometric variables (Scholz & Eislöffel 2004) or low-mass objects (Tej et al. 2002; Sherry et al. 2004) is still prevalent. The summit of our relative poor knowledge of the stellar and high-mass brown-dwarf population in the cluster is maybe the case of SE 70. Scholz & Eislöffel (2004) detected it at 4.6 arcsec to a previously known $\sim 5 M_{\text{Jup}}$ -mass candidate planetary-mass object, S Ori 68 (Zapatero Osorio et al. 2000). Caballero et al. (2006a) have recently confirmed the membership in the cluster of SE 70, which is an X-ray-emitter brown dwarf² and may form together with S Ori 68 the widest planetary system ever found (1700 ± 300 AU). The recent discovery of at least two spatially coincident stellar populations in the area makes current and future studies more complex. The superimposed stellar components, firstly showed by Jeffries et al. (2006) and confirmed by Sacco et al. (2006), are kinematically separated in radial velocity and have different mean ages. Furthermore, σ Orionis itself might not be a member of any of the known Ori OB1 subgroups, but be part of a new one. For example, the simple fact of the Horsehead Nebula in the Ori OB 1b Association being a back-illuminated absorption nebula may suggest that it is at a different heliocentric distance than σ Orionis.

A complete study of the stellar population in the σ Orionis cluster is absolutely needed to complement current and future deep searches of brown dwarfs and planetary-mass objects in the area (e.g. to compare spatial distributions, disc frequencies or slopes of the mass spectrum at different mass intervals and separations to nearby massive stars). In this paper, I will describe an optical-near infrared search for the most massive stars in the area of σ Orionis, supported by astrometric, X-ray and mid-infrared measurements. The search and characterization of intermediate- and low-mass stars in the cluster will be presented in a forthcoming paper.

2. Analysis

A cross match between the Tycho-2 and 2MASS catalogues will lead to the identification of the brightest stars (in the optical) of the Ori OB 1b Association in the direction of the σ Orionis cluster. The collection of additional data from different archives related to mid-infrared and X-ray space missions and from the literature will provide information on the nature and age of the stars.

2.1. The Tycho-2/2Mass correlation

For the present work, I have used the web-based sky atlas Aladin (v3.600; Bonnarel et al. 2000; Fernique et al. 2004). I have loaded all the sources catalogued in the Tycho-2 and the 2MASS Point Source catalogues (Høg et al. 2000 and Cutri et al. 2003, respectively) in a circle of 30 arcmin radius centred in σ Orionis A+B+IRS1 (the close triple OB system formed by σ Ori A, σ Ori B and σ Ori IRS1 that gives the name to the cluster). In the studied area of ~ 2830 arcmin², there are 41 Tycho-2 and about 50 000 2MASS catalogued sources. Previous studies in the

² Burningham et al. (2005) also measured the equivalent width of the Na I $\lambda\lambda 8183, 8195$ Å doublet and the radial velocity of SE 70, which are very similar to those of other confirmed young brown dwarfs in the σ Orionis cluster.




Fig. 1. Inverse-colour DSS-1 blue-band (photographic B_I) image centred in σ Ori A (marked with an arrow). Size is $1.5 \times 1.5 \text{ deg}^2$. North is up and East is left. Sources in the Tycho-1 and Tycho-2 are marked with (red) crosses and (blue) squares, respectively (in colour in the electronic version). East and north-east of the field of view are dominated by the star-forming region complex formed by Alnitak, the Flame Nebula (NGC 2024), the Horsehead Nebula and the IC 434 H II region.

region pointed out that 30 arcmin ($\sim 3 \text{ pc}$ at the distance of σ Orionis) is a rough but valid approximation to the cluster radius (Sherry et al. 2004; Béjar et al. 2004a). Many bright stars out of this radius may be related to the youngest star-forming regions to the east of the Ori OB 1b Association or to the Alnilam cluster to the west (see e.g. Wolk 1996). The B1.5V-type star HD 37744 and the Horsehead Nebula are in the vicinity of the studied region.

The positional cross match in the area with the Aladin Cross match tool between the Tycho-2 and 2MASS catalogues provided a list of 41 stellar sources. Therefore, the 2MASS near-infrared counterparts of all the Tycho-2 sources in the area were identified. I used the default cross-match threshold of 4 arcsec. Except in one case, the agreement between the Tycho-2 and 2MASS coordinates was better than 0.5 arcsec (and better than 0.2 arcsec in 78 % of the cases). The difference between the Tycho-2 and 2MASS coordinates of the outlier was 2.4 arcsec. This source (TYC 4770 1432 1), with a magnitude in the reddest Tycho band $V_T = 12.3 \pm 0.2 \text{ mag}$, is one of the faintest stars in the sample. The deviation may be ascribed to the low signal-to-noise ratio in the Tycho catalogue, which 90 %-completeness magnitude in the V band is about 11.5 mag (Høg et al. 2000).

Figure 1 shows a $\sim 1.5 \times 1.5 \text{ deg}^2$ blue digitized photographic plate, covering the region of 30 arcmin centred in σ Ori A and adjacent areas. The 41 correlated stars are marked in the view window with squares. For each star, it is available photometry in five bands ($B_T V_T JHK_s$) and accurate proper motion measurements. In the upper parts of Tables A.1 and A.2, I provide their names and the photometric and astrometric data, respec-

Table 2. Mid-infrared sources among the stars in the Tycho-2/2MASS correlation.

Name	F_{12} [Jy]	F_{25} [Jy]	F_{60} [Jy]	F_{100} [Jy]
σ Ori A+B+IRS1 ^a	4.5 ± 0.2	15 ± 2	15 ± 2	15 ± 2
HD 37699	0.29:	0.45 ± 0.09	1.5 ± 0.2	4.9:
HD 294268	0.87 ± 0.11	2.74 ± 0.16	1.50 ± 0.10	22:

^a Re-measured IRAS photometry (Oliveira & van Loon 2004).

tively³. The stars in the Tycho-1 catalogue (ESA 1997), superseded in most applications by Tycho-2, are marked with crosses in Figure 1. There are four additional stars that are in Tycho-1 and not in Tycho-2.

2.2. Additional data

2.2.1. IRAS space telescope

Three out of the 41 correlated stars possess an unambiguous source of the IRAS catalogue of Point Sources Version 2.0 (IPAC 1986) at less than 40 arcsec. Their names and average non-colour corrected flux densities, F_ν , for 12, 25, 60 and 100 μm are given in Table 2. The three stars have been investigated as mid-infrared sources with probable discs (σ Ori A+B+IRS1 – van Loon & Oliveira 2004; HD 294268 – García-Lario et al. 1990, 1997 and Torres et al. 1995; HD 37699 – Oganesyan et al. 1995).

In the studied area there are only four additional IRAS unambiguous sources associated to fainter stars and their surrounding envelopes. They are TX Ori, V510 Ori, IRAS 05358–0238 and IRAS 05352–0227. None of them are in the Tycho-2 catalogue. However, the correlated star HD 294269 has an unreliable detection at only one passband, 60 μm , and will not be considered.

2.2.2. ROSAT, XMM-Newton and Chandra space telescopes

At least 13 stars of the sample are X-ray emitters. They were found in the following catalogues: ROSAT Source Catalog of Pointed Observations with the High Resolution Imager, 1RXH (ROSAT 2000a); WGACAT version of the ROSAT PSPC Catalogue, WGACAT (White et al. 2000); Second ROSAT Source Catalog of Pointed Observations, 2RXP (ROSAT 2000b).

All the X-ray emitters, except σ Orionis E and the faint star 4771–0950 (Wolk 1996), are present in the WGACAT-ROSAT catalogue. Eight were previously detected in the spatial and spectral analysis of a full EPIC field in the XMM-Newton observations of the σ Orionis cluster by Franciosini et al. (2006). They reported deviations between the coordinates of the optical and the X-ray sources of less than 1 arcsec for all the bright stars in their sample. Additionally, HD 37699 and HD 294268, to the northeast of the area, were also detected in the X-ray observations of the Orion OB 1b Association with the ASCA mission by Nakano et al. (1999). Voges et al. (1999) considered GSC 04771–00962 and GSC 04771–00961 as possible HST Guide Star Catalogue optical counterparts of the X-ray ROSAT source 1RXS J053756.8–023857, which is in reality ascribed to the

³ Tables A.1 and A.2 are only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/>.

Table 3. X-ray sources among the stars in the Tycho-2/2Mass correlation^a.

Name	Nr. 1RXH	WGACAT [10^{-3} s^{-1}]	XMM [10^{-3} s^{-1}]
σ Ori A+B+IRS1	129	330±80	440±2
σ Ori E	77	–	199.4±1.7
HD 37699	4	90±20	×
HD 37525 AB	129	19.6±1.3	–
HD 294272 A+B	3	3.6±0.6	5.9±0.4
HD 37564	4	24±5	9.9±0.5
V1147 Ori	×	70±40	×
HD 294268	×	3.50±0.11	×
RX J0539.6–0242 AB	142	60±20	63±2
HD 294298	×	60±8	×
4771–0950	1	–	0.67±0.18
2E 0535.4–0241	160	53±17	189±4

^a Name of optical counterpart, number of events in 1RXH-ROSAT, count rate (s^{-1}) in WGACAT-ROSAT and count rate (s^{-1}) in the XMM-Newton observations by Franciosini et al. (2006). The ROSAT and XMM-Newton missions are not sensitive to the same wavelength interval. The symbol “×” denotes that the target is not in the area studied by the X-ray mission or catalogue.

bright young star 2E 0535.4–0241. Table 3 shows the X-ray emitters in the sample, the count rates and the number of events measured by different instruments and space missions.

The high background produced by the extreme X-ray emitter σ Ori A+B+IRS1 did not allow to recognise σ Ori E as an individual WGACAT source. As well, the small angular separation between HD 294272 A and B (~ 8 arcsec) and their similarity in brightness did not permit to discriminate which is the real X-ray emitter from ROSAT and XMM-Newton data. However, both σ Ori A+B+IRS1 and σ Ori E and HD 294272 A and B systems are resolved in the high-spatial-resolution X-ray images from the *Chandra* Space Telescope archive obtained by Adams et al. (2004) with HRI-C.

The null detection at high energies of the rest of the targets, except in the case of StHA 50, may be due to the large separation to σ Ori A, where X-ray surveys are usually centred.

2.2.3. Optical spectroscopy

From my searches in the literature, 28 stars have a spectral type determination. Their spectral types are provided in the last column of Table A.2. Out of them, 13 stars (14 if the Be-type star StHA 50 is also included) have OBA spectral types. Other five stars have F types, and the rest are G- and K-type stars. The spectral type determinations have been borrowed from the following works: Schild & Chaffee (1971), Warren & Hesser (1978), Guetter (1981), Downes & Keyes (1988), Gray & Corbally (1993), Nesterov et al. (1995), Wolk (1996), Catalano & Renson (1998), Alcalá et al. (2000), Gregorio-Hetem & Hetem (2000) and Caballero (2006).

There is information about the lithium abundance for seven stars. In six cases, the equivalent width of the $\text{Li I } \lambda 6707.8 \text{ \AA}$ suggests that the lithium abundance is primordial ($\log \epsilon(\text{Li}) \sim 3.0$), while there is an upper limit of 1.0 dex for the $\log \epsilon(\text{Li})$ of the G0-type star HD 294269 (Cunha, Smith & Lambert 1995; Torres et al. 1995; Alcalá et al. 1996, 2000; Caballero 2006). See in

Table 4. Very young stars in the Tycho-2/2MASS correlation.

Name	OB	X-ray	Li I	H α	mIR	var.
σ Ori A+B+IRS1	Y	Y			Y	
σ Ori E	Y	Y				Y
HD 37699	Y	Y			Y	
HD 294271	Y					
HD 37525 AB	Y	Y				
HD 294272 A	Y	Y				
HD 294272 B	Y	Y				
HD 37333	Y					
HD 37564		Y				
V1147 Ori	Y	Y				Y
HD 37686	Y					
HD 37545	Y					
HD 294275	Y					
HD 294268		Y	Y	Y	Y	Y?
RX J0539.6–0242 AB		Y	Y	Y		
HD 294279			Y			
StHA 50	Y?			Y		
HD 294298		Y	Y			
4771–0950		Y	Y			
2E 0535.4–0241		Y				
SO120532			Y			

Cunha et al. (1995) a careful study of the lithium abundance of late F and early G stars in Orion and how very young ages can be derived for stars with effective temperatures between ~ 5200 and ~ 6400 K.

There are three stars with H α in emission: RX J0539.6–0242 AB ($\text{EW}(\text{H}\alpha) = -2.38 \text{ \AA}$; Alcalá et al. 1996), StHA 50 (Stephenson 1986; Downes & Keyes 1988) and HD 294268 ($\text{EW}(\text{H}\alpha) = -2 \text{ \AA}$; Torres et al. 1995). As well, seven stars display spectroscopic peculiarities: V1147 Ori (abnormally strong silicon and europium lines; Catalano & Renson 1998), HD 37333 (mild Si II; Gray & Corbally 1993), HD 37699, HD 37686 and HD 294301 (broad “nebulous”-absorption due to spinning; Neubauer 1943, Warren & Hesser 1978 and Gray & Corbally 1993, respectively), HD 37525 AB (helium weak; Nissen 1976) and σ Ori E (helium rich and magnetic strong; Greenstein & Wallerstein 1958). The stars HD 294298 and RX J0539.6–0242 AB have also large values of $v \sin i$ if compared with field solar-like stars (55 and 150 km s^{-1} , respectively; Cunha et al. 1995 and Alcalá et al. 1996). Finally, Hunger et al. (1989) measured a $v \sin i$ of $160 \pm 20 \text{ km s}^{-1}$ in σ Ori E.

2.3. Target classification

2.3.1. Young stars

The primordial abundance of lithium (“Li I”), the line of H α in appreciable emission in solar-type stars or earlier (“H α ”), the mid-infrared flux excess in the *IRAS* pass-bands due to a disc (“mIR”), the very early spectral types (O, B and early A; “OB”), the X-ray emission (strong if a solar-type star; “X”), the high-amplitude photometric variability (“var.”) and the fast rotation are some known youth features (age $\lesssim 10$ Ma). These signatures are common in T Tauri stars and their analogous at intermediate masses, the Herbig Ae/Be stars (with masses in the interval $1.5 M_{\odot} < M < 10 M_{\odot}$ and spectral types earlier than F; Hillenbrand et al. 1992). Out of the 41 stars of the Tycho-2




Fig. 2. Proper motions of the 41 Tycho-2/2MASS-correlation stars. Code: (red) filled stars, “★”, young stars; (blue) crosses, “×”, field stars with proper motion different from the mean proper motion of the Ori OB 1b Association; (blue) pluses, “+”, probable K- or M-type foreground stars; (red) open circles, “○”, possible new Ori OB 1b Association members. The dotted and solid big circles (ellipses with the axes scale) denote the Brown et al. (1999) and this paper astrometric selection criteria, respectively. The object marked with an asterisk, “*”, is HD 294278, which is a K-type foreground star with a different proper motion (“×” and “+” simultaneously).

sample, 21 have at least one youth feature. In Table 4, I provide the list of the very young stars. I will consider that they are bona-fide members of the Ori OB 1b Association, to which the σ Orionis cluster belongs. Some of the stars display several youth features at a same time, as HD 294268, which is a mid-infrared source with moderate X-ray and H α emissions and lithium in absorption. It is also a probable photometric variable star. Warren & Hesser (1978) suggested the membership to the Ori OB 1b Association of 14 stars among the subsample of young stars in Table 4.

2.3.2. Stars with large tangential velocity

The tabulated mean proper motion of all the subgroups of the Ori OB1 Association is $\mu_\alpha \cos \delta = +0.44 \text{ mas s}^{-1}$, $\mu_\delta = -0.65 \text{ mas s}^{-1}$ (Brown et al. 1999). de Zeeuw et al. (1999) showed that the tangential velocities of 96 % of the Ori OB1 photometric candidates identified by Brown et al. (1994) deviated less than 5 mas s^{-1} from that mean proper motion. However, in the work presented here, there is a large fraction of young stars that display features of extreme youth and their tangential velocity do not satisfy the de Zeeuw et al. (1999) proper-motion criterion. One extreme case is that of the F7-type X-ray-emitter star 4771–0950, with $p\text{EW}(\text{Li I}) = +0.07 \pm 0.01 \text{ \AA}$ (Caballero 2006), which proper motion deviates in almost 10 mas s^{-1} from the mean proper motion of the Ori OB1 Association. I have used the value of 10 mas s^{-1} to separate young stars and candidate Association members from probable foreground stars with larger tangential velocities. This wider “error circle” is not caused by larger errors in the proper motions of the faintest members, but due to a genuine tangential velocity dispersion (see Section 3.3). Figure 2 illustrates the astrometric study, showing the selection criteria by Brown et al.

Table 5. Stars in the Tycho-2/2MASS correlation with proper motion different from that of the Ori OB1 Association in more than 10 mas s^{-1} .

Name	$\mu_\alpha \cos \delta \pm \delta\mu_\alpha \cos \delta$ [mas a ⁻¹]	$\mu_\delta \pm \delta\mu_\delta$ [mas a ⁻¹]	Sp. type
HD 294307 ^a	+0.1±1.4	−16.8±1.4	F8
GSC 04771–00621	+7.0±1.6	−11.1±1.6	
HD 294278	+10.9±1.4	+2.6±1.4	K2
HD 294270 ^b	+16.8±1.9	−23.4±1.8	G0
HD 294269 ^c	+43.5±1.8	−16.1±1.8	G0
HD 294274	−8.1±3.2	−19.4±3.5	G0
TYC 4771 720 1	−20.1±3.9	−16.6±4.2	
TYC 4771 661 1	+16.6±4.3	−20.1±4.7	
TYC 4770 1432 1	−30.9±4.5	+0.1±5.0	
TYC 4770 924 1	+1.8±4.7	−43.4±5.1	

^a Non member of the Ori OB 1b Association in Warren & Hesser (1978).

^b Uncertain association membership status in Warren & Hesser (1978).

^c No detection of Li I in absorption and $V_r = +72.8 \text{ km s}^{-1}$ in Cunha et al. (1995).




Fig. 3. Same as Figure 2 but for the V_T vs. $V_T - K_s$ colour-magnitude diagram.

(1999) and the one used here. The ten stars with tangential velocity different from that of the Ori OB1 Association in more than 10 mas s^{-1} are shown in Table 5 (they could be at up to 3 deg from σ Ori AB in 1 Ma).

2.3.3. K- and M-type foreground stars

In the V_T vs. $V_T - K_s$ colour-magnitude diagram shown in Figure 3, there are two well-differentiated regions separated at $V_T \sim 9.5 \text{ mag}$. On the one hand, the 13 stars brighter than this value are bluer than $V_T - K_s = 1.0 \text{ mag}$, display youth features and, hence, verify the astrometric criterion. On the other hand, out of the 28 stars fainter than $V_T \sim 9.5 \text{ mag}$, ten are the proper-motion outliers shown in Table 5. There are also seven stars, shown in

Table 6. Probable K- and M-type foreground stars in the Tycho-2/2MASS correlation.

Name	Sp. type
HD 294277 ^a	K2
HD 294278 ^b	K2
HD 294280 ^a	K5
TYC 4770 1018 1	
TYC 4771 1012 1	
TYC 4771 934 1	
TYC 4771 1468 1	

^a They were found by the Midcourse Space Experiment satellite at 8.3 μ m (Kraemer et al. 2003).

^b It does not satisfy the astrometric selection criterion either.

Table 7. Possible new Ori OB 1b association member stars in the Tycho-2/2MASS correlation.

Name	Sp. type
HD 294301 ^a	F2V(n)
GSC 04771-00962	
TYC 4770 1261 1	
TYC 4770 1129 1	

^a Uncertain association membership status in Warren & Hesser (1978).

Table 6, with $V_T - K_s$ colours redder than ~ 3.0 mag, which are located far from the photometric cluster sequence defined by the young stars. Their $V_T - K_s$ colours and V_T apparent magnitudes are typical of early/intermediate K dwarfs in foreground. Furthermore, Nesterov et al. (1995) determined K spectral type for the three brightest stars. Given their brightness (with J magnitudes brighter than 9.5 mag), if they were association members, then they should be extraordinary reddened stars showing the astrophysical properties of Class I objects. None of them shows X-ray or mid-infrared emissions detectable by present surveys, nor any spectroscopic youth feature. On the contrary, the three young members of the Ori OB 1b Association with $V_T - K_s$ colours between 2.0 and 2.5 mag and of similar brightness in the V_T band to the stars in Table 6 are among the strongest X-ray emitters in the σ Orionis cluster (RX J0539.6-0242 AB, 2E 0535.4-0241 and HD 294298).

2.3.4. Possible new Ori OB 1b Association members

There are four stars that have not been classified as young stars, proper-motion outliers or foreground K-type stars, and that follow the photometric sequence of the normal young stars. They are tabulated in Table 7. There is available spectroscopic information only for the fast-rotating F2-type star HD 294301 (Gray & Corbally 1993). The star GSC 04771-00962 might be related to an unidentified X-ray event in the Second ROSAT PSPC Catalog, at 0.7 arcmin to the north of the star. I will consider the four stars as possible new Ori OB 1b Association members.

2.4. Binary systems

Among the studied targets, there are at least four close multiple systems. They are σ Ori A+B+IRS1, HD 37525 AB, HD 294272 A and B and the spectroscopic binary RX J0539.6-0242 AB. Both Tycho-2 and 2MASS observations were able to resolve the HD 294272 system. It is not the case, however, of the other three stars. The secondary at 0.45 ± 0.4 arcsec of HD 37525 A is more than 0.5 mag fainter in the H band than the primary, so the contribution of the B component to the total flux at optical wavelengths may not be appreciable. The same occurs, in an extreme case, with σ Ori IRS1, which is more than 5 mag fainter than σ Ori AB in the H band (Caballero 2005, 2006). However, the optical flux of σ Ori B is only three times less than that of σ Ori A. Although the short separation between both components (of 0.250 arcsec in J1991.25), it is possible to estimate the optical magnitudes of σ Ori A and B from Hipparcos data. The combined H_P magnitude of σ Ori AB is 3.681 ± 0.016 mag, while the magnitude difference between them, ΔH_P , is 1.21 ± 0.05 mag. Using the $B - H_P$ and $H_P - V$ colours of blue Hipparcos stars as a reference, I have estimated that the Johnson B - and V -band magnitudes of σ Ori A and B are 3.85 and 4.10 mag, and 5.18 and 5.26 mag, respectively, with errors of ~ 0.05 mag. I have used the relationships $\Delta m_{A+B} = m_B - m_A$ and $m_A = m_{A+B} + 2.5 \log \left(1 + 10^{-\frac{\Delta m_{A+B}}{2.5}} \right)$. The V_A and V_B estimations nicely agree with V -band adaptive optics observations of σ Ori AB by ten Brummelaar et al. (2000). The spectroscopic binary RX J0539.6-0242 AB cannot be resolved by imaging with current technology. Tovmassian et al. (1991) also proposed a suspected binary among the correlated stars, HD 37564, but it has not been confirmed.

2.5. Missing stars

2.5.1. Young stars in the Tycho-1 catalogue

The Tycho-2 catalogue is not complete. As mentioned before, there are four stars in Tycho-1 that are not in Tycho-2. Their basic properties (photometry, astrometry, spectral types) are tabulated in the bottom parts of Tables A.1 and A.2. The stars are brighter than or as bright as the Tycho V_T completeness magnitude. Furthermore, one of them is σ Ori D, which is one of the brightest stars in the σ Orionis cluster. The four stars roughly follow the cluster sequence in the V_T vs. $V_T - K_s$ colour-magnitude diagram. Their proper motion measurements are affected by effects external to the Hipparcos satellite (like the glare of σ Ori A+B+IRS1 in the case of σ Ori D or the background emission of the Horsehead Nebula in the case of HD 294297). Errors in their proper motions in Tycho-1 are larger than 5 mas a^{-1} and up to 30 mas a^{-1} , and Tycho-2 not even tabulates them. Table A.2 shows the tangential velocities of the four stars provided by Høg et al. (1998), which matches at the level of a few mas a^{-1} with the USNO-B1 and NOMAD1 catalogues (Monet et al. 2003 and Zacharias et al. 2004, respectively).

Out of the four missing stars, three display youth features (Cunha et al. 1995; Sanz-Forcada et al. 2004; Caballero 2005, 2006). The young missing stars are shown in Table 8. Very interestingly, HD 294297 has both a primordial lithium abundance and a proper motion that clearly deviates from the rest of the Ori OB 1b Association members. Its tangential velocity is $(\mu_\alpha \cos \delta, \mu_\delta) = (+22.7 \pm 2.6, -20.5 \pm 1.7) \text{ mas a}^{-1}$ from Høg et al. (1998) and $(+22.1 \pm 1.6, -23.9 \pm 1.4) \text{ mas a}^{-1}$ from the NOMAD1 catalogue. The remaining missing target, the G0-type star HD 294276, does not display any youth feature and has the largest

Table 8. Young stars in the Tycho-1 catalogue and not in the Tycho-2 catalogue.

Name	OB	X-ray	Li I	Sp. type
σ Ori D ^a	Y	Y		B2V
HD 294273	Y			A3
HD 294297			Y	F6–8

^a $v \sin i = 180 \pm 20 \text{ km s}^{-1}$ (Hunger et al. 1989).

tangential velocity among the stars studied in this work, so it is likely a foreground solar-like star.

2.5.2. σ Ori C

The Hipparcos mission was mostly sensitive to the blue optical wavelengths. There are no stars brighter than $J = 9.0 \text{ mag}$ in the 2MASS catalogue and in the surveyed area without optical counterpart in the Tycho catalogues. However, there are 12 2MASS stars with magnitudes $9.0 < J \leq 10.0 \text{ mag}$ not found in Tycho-1 or 2. Among them, there are four foreground K- and M-type stars (based on mid-resolution optical spectroscopy and/or proper motion; Caballero 2006), five cluster members and three stars of unknown nature. Out of the five young stars, the brightest one is the A2-type star σ Ori C, which is about 2.7 mag brighter in the V band than the 90%-completeness magnitude of the Tycho-2 catalogue. Its absence in the Tycho catalogues may be due to the same reason as σ Ori D in the optical and σ Ori E in X rays: the glare of σ Ori AB. I will “recover” σ Ori C as one of the brightest σ Orionis cluster members.

3. Discussion

To sum up, out of the 41 stars in the Tycho-2 catalogue at angular separations less than 30 arcmin of σ Ori A, 21 stars have youth features according to my search in catalogues and in the literature. I have also computed the B - and V -band magnitudes of σ Ori A and B separately and considered four additional bright young stars (three from the Tycho-1 catalogue plus σ Ori C) and four new candidate Ori OB 1b Association members. This makes a list of 26 bright young stars and four candidate young stars in the studied area. Out of them, nine are in multiple systems: σ Ori A, B, C, D and E, HD 294272 A and B, HD 37525 AB and RX J0539.6–0242 AB. It indicates that the minimum frequency of multiplicity of the brightest stars is 35 % (the low-mass stars and substellar objects are not taken into account). Next, I will characterize the sample of young stars.

3.1. Individual masses and the V vs. $B - V$ diagram

For each of the 30 bright young stars and candidates, I have derived their masses from the Johnson B - and V -band and theoretical stellar evolution isochrones from the Geneva database (Lejeune & Schaerer 2001). Johnson magnitudes were derived from Tycho B_T and V_T magnitudes using the relations $V = V_T - 0.090(B_T - V_T)$ and $B - V = 0.850(B_T - V_T)$, which are valid in the colour interval $-0.2 \text{ mag} < B_T - V_T < 1.8 \text{ mag}$ (ESA 1997). For σ Ori A and B, I have used the B and V magnitudes derived in Section 2.4, while for σ Ori C, I have used the photometry from Greenstein & Wallerstein (1958).




Fig. 4. V vs. $B - V$ colour-magnitude diagram of the young stars in the Tycho-2/2MASS correlation (filled stars), the four new probable Ori OB 1b Association members (open circles), σ Ori B, C and the three young stars in the Tycho-1 recovery (filled circles). Solid, dashed and dot-dashed lines (in overlapping) denote the 1-, 3- and 10-Ma isochrones of the Geneva group at the most probable heliocentric cluster distance, and the dotted lines indicate the 3-Ma isochrones at the minimum (300 pc) and maximum (430 pc) possible cluster distances. Several target are labeled.

The V vs. $B - V$ colour-magnitude diagram depicted in Figure 4 illustrates the mass derivation from the models. In particular, I have used the basic grid for solar metallicity $Z = 0.02$ (Schaller et al. 1992) and conservative possible intervals in the age (3^{+7}_{-2} Ma) and in the heliocentric distance ($360^{+70}_{-60} \text{ pc}$) of σ Orionis. The most probable masses for each target and for each age-distance pair was computed from the mass- M_V relationship given by the models. The masses with their uncertainties for each target are shown in Table 9. The edge of the conservative error bars correspond to the extreme cases of a younger closer cluster (1 Ma, 300 pc) and for an older farther cluster (10 Ma, 430 pc), which in contrast are close to be ruled out by the isochronal matches to low-mass stars. The extended grid with high mass loss for massive stars (Meynet et al. 1994) provided identical results, except for σ Ori A, which derived mass slightly varied within the uncertainties of the age and heliocentric distance of the cluster. The masses of HD 37525 AB and RX J0539.6–0242 AB were not corrected from binarity.

There are three bright stars ($V < 9.5 \text{ mag}$) that are 0.1–0.2 mag redder than the isochrones and the sequence defined by the other cluster members in the colour-magnitude diagram. They are σ Ori B (which atmosphere could be suffering the effects of the O9.5V primary at only $\sim 90 \text{ AU}$), σ Ori E (a helium rich star – i.e. the used models may be not applicable) and HD 37564 (which may have a disc – see Section 3.2). The masses derived in this work for these objects must be cautiously treated. The rest of the bright stars also lie slightly to the red of the isochrones in the colour-magnitude diagram in Figure 4. Since in the high-mass domain of the colour-magnitude diagram in Fig. 4 there is almost no sensitivity to the age of the cluster, this redshift favours the nearest possible heliocentric distances to σ Orionis. It implies thus that the cluster could be older than usually assumed. This fact of the greatest importance for the

Table 9. Masses of the young stars.

Name	Mass [$M_{\odot}$]	Remarks ^a
σ Ori A	18 ± 6	XX, (mIR)
σ Ori B	12 ± 3	XX
σ Ori E	$7.4^{+1.5}_{-1.4}$	He rich, XX
σ Ori D	$6.8^{+1.8}_{-1.2}$	
HD 37699	$4.4^{+0.8}_{-0.7}$	mIR, XX
HD 294271	$3.9^{+0.8}_{-0.6}$	
HD 37525 AB	$3.6^{+0.7}_{-0.6}$	binary, XX
HD 294272 A	3.2 ± 0.5	
HD 294272 B	3.0 ± 0.5	
HD 37333	$3.0^{+0.6}_{-0.4}$	
HD 37564	$3.1^{+0.6}_{-0.4}$	
σ Ori C	2.7 ± 0.4	
V1147 Ori	$3.5^{+0.4}_{-0.3}$	XX
HD 37686	$2.3^{+0.4}_{-0.3}$	
HD 37545	$2.2^{+0.3}_{-0.1}$	
HD 294275	$2.1^{+0.3}_{-0.1}$	
HD 294297	$1.7^{+0.2}_{-0.1}$	μ
HD 294268	$1.6^{+0.2}_{-0.1}$	mIR
RX J0539.6–0242 AB	1.8 ± 0.1	SB, XX
HD 294279	1.6 ± 0.1	
HD 294273	1.5 ± 0.1	
StHA 50	1.4 ± 0.1	
HD 294301	1.5 ± 0.1	prob.
GSC 04771–00962	1.4 ± 0.1	prob.
HD 294298	1.5 ± 0.1	XX
4771–0950	1.3 ± 0.1	
2E 0535.4–0241	1.5 ± 0.1	XX
TYC 4770 1261 1	1.3 ± 0.1	prob.
TYC 4770 1129 1	1.3 ± 0.1	prob.
SO120532	1.2 ± 0.1	


^a Remarks: “mIR”: IRAS source; “XX”: moderate or strong X-ray emitter; “prob.”: probable association member star without detected youth features. The rest are described in the text.

derivation of the mass function in the cluster substellar domain down to planetary masses (Béjar et al. 2001; González-García et al. 2006; Caballero et al. 2007a), since the theoretical masses from the models strongly depend on the age. An age older than expected for σ Orionis, at about 10 Ma, would lead dozens currently considered brown dwarfs to be very low-mass stars, and several planetary-mass objects to be really brown dwarfs.

3.2. Disc harbours and X-ray emitters

Some of the young stars in Table 9 display characteristics that are present in T Tauri and Herbig Ae/Be stars. In the standard scenario, these stars possess discs, which are responsible of the flux excess in the mid and/or near infrared. The in-falling material from the disc onto the stellar surface produces the X-ray emission and the strong broad H α emission. The columns of the accreting mass hitting the atmosphere can produce photometric variability and flux excess in the blue optical as well (Bertout et al. 1989; Appenzeller & Mundt 1989; Königl 1991; Hillenbrand et al. 1992).

Among the investigated stars, three were detected by the IRAS space telescope (Table 2 in Section 2.2.1). I have built their spectral energy distributions (SEDs) from the blue optical to 100 μ m ((U)BVRIJHK_s(L'[8.6])[12][25][60][100]), using

**Fig. 5.** Spectral energy distributions of σ Ori A+B+IRS1 (top), HD 37699 (middle) and HD 294268 (bottom). Vertical scales are different.**Table 10.** Young stars with probable discs^a.

Name	Sp. type	Remarks
HD 37699	B5Vn	IRAS source
HD 37564	A8V:	red $B - V$
HD 294268	F5	IRAS source
RX J0539.6–0242 AB	G5–K0	$V_T - K_s > 2.0$ mag
StHA 50	Be	blue $B - V$
HD 294298	G0:	$V_T - K_s > 2.0$ mag
2E 0535.4–0241		$V_T - K_s > 2.0$ mag

^a Discarding σ Ori IRS1.

^b Weak-line T Tauri binary (Alcalá et al. 1996).

the data presented here and from Lee (1968) and van Loon & Oliveira (2004). There is high-resolution (sub-arcsec) imaging for only the σ Ori A+B+IRS1 system. The mid-infrared source σ Ori IRS1, at 3.3 arcsec northeast to the OB system σ Ori AB, has been resolved at 6–20 μ m (van Loon & Oliveira 2003), in X rays (Sanz-Forcada et al. 2004), in the J and H bands (Caballero 2005) and in the optical (Rebolo et al. in prep.). It is the responsible of the IRAS emission coming from the triple system. I propose that σ Ori IRS1 is a low-mass K:-type star surrounded by a thick envelope that loses its outer atmospheric layers due to the strong ultraviolet radiation field and stellar wind coming from the O9.5V+B0.5V system at only ~ 1000 AU (see also van Loon & Oliveira 2003). The other two stars with IRAS emission, HD 37699 and HD 294268, have masses of $4.4^{+0.8}_{-0.7}$ and $1.6^{+0.2}_{-0.1} M_{\odot}$, respectively, which make them to be suitable candidate Herbig Ae/Be stars. Besides, according to Bertout (1989), the more massive T Tauri stars rotate significantly faster than less massive ones. The large $v \sin i$ derived from the “nebulous” spectral type of HD 37699 (Neubauer 1943) points out that this relation may be extrapolated to Ae/Be stars. The disc surrounding HD 37699 seems also to be cooler than those of σ Ori IRS1 and HD 294268, since the flux density at 60 μ m is larger than at 25 μ m, to the contrary to what happens in the other two. The large uncertainties in the IRAS fluxes prevent to derive additional properties of the discs. The abrupt step in the SED of HD 294268




Fig. 6. $J - H$ vs. $H - K_s$ colour-colour diagram of the brightest stars of the σ Orionis cluster. Stars with and without disc are marked with filled stars and open circles, respectively. The error bars of σ Ori AB have not been drawn for clarity.

between the $BVRI$ -band data (Lee 1968) and the 2MASS JHK_s -band data (Cutri et al. 2003) suggests a secular large-amplitude photometric variability.

There are five additional candidate star systems with probable discs. They are shown, together with HD 37699 and HD 294268, in Table 10. Three of them are the strong X-ray emitters with colours $V_T - K_s > 2.0$ mag (Section 2.3.3). The remaining probable disc harbours are HD 37564 and StHA 50. The former is an A8-type star with moderate X-ray emission and relatively red $B - V$ and $V_T - K_s$ colours. StHA 50, for which there is only a few data available (it is a Be-type emission-line object with $V \approx 11.4$ mag – Stephenson 1986; Downes & Keyes 1988), deviates at the level of 3σ to the blue in the V vs. $B - V$ colour-magnitude diagram in Figure 4 respect to the sequence defined by the other cluster members. IRAC *Spitzer* Space Telescope data between 4.5 and $8.0\mu\text{m}$ supports the hypothesis of discs surrounding both HD 37564 and StHA 50 (Caballero et al. in prep.). As a matter of fact, both of them and HD 37699, the bluest star with a disc in σ Orionis, have no appreciable near-infrared flux excess in the $J - H$ vs. $H - K_s$ colour-colour diagram shown in Fig. 6. Their discs are probably cooler than those surrounding the remaining stars in Table 10.

The frequency of discs surrounding young stars in the magnitude interval $7.5\text{ mag} \lesssim V \lesssim 11.0\text{ mag}$ and in the direction of σ Ori is $32 \pm 14\%$. This value is of the order of other disc frequencies measured in the area in different mass intervals (compiled in Table 11). The frequencies derived from the detection of flux excess red-wards $5\mu\text{m}$ are in general larger than the disc frequencies based in only JHK_s photometry. Since the various disc frequencies quoted have been assessed heterogeneously with different techniques, I do not claim any evidence for a mass-dependence of the disc frequency in σ Orionis.

The strongest X-ray emitters are σ Ori A+B+IRS1 and σ Ori E. The σ Ori A+B+IRS1 system was resolved in the AB and IRS1 components with HRI-C/*Chandra* Space Telescope observations. The bulk of the X-ray flux comes from the OB pair (Sanz-Forcada et al. 2004). Other moderate emitters are HD 37699, HD 37525 AB, V1147 Ori and the three stars with

Table 11. Frequency of discs in different mass intervals in the Ori OB 1b Association and in the σ Orionis cluster.

Frequency (%)	Reference	Method ^a
17^{+18}_{-17}	Jawawardhana et al. (2003)	Ja03
5–12	Barrado y Navascués et al. (2003)	ByN03
46–54	Oliveira et al. (2004)	Ol04
~25	Caballero et al. (2004)	Ca04
5–7	Scholz & Eislöffel (2004)	SE04
17.4 ± 3.6	Hernández et al. (2005)	He05
10 ± 5	Kenyon et al. (2005)	Ke05
46^{+16}_{-13}	Caballero (2005)	Ca05
33 ± 6	Oliveira et al. (2006)	Ol06
33 ± 4	Caballero et al. (2006b)	Ca06b
47 ± 14	Caballero et al. (2007a)	Ca07a
30 ± 20	Caballero et al. (2007b)	Ca07b
32 ± 14	This work	

^a (Ja03): very low-mass stars in σ Orionis with L' excess; (ByN03): intermediate- and low-mass stars and high-mass brown dwarfs in σ Orionis with K_s excess (minimum frequency); (Ol04): low-mass stars and high-mass brown dwarfs in σ Orionis with $K_s L'$ excess; (Ca04): brown dwarfs in σ Orionis with K_s excess, $H\alpha$ emission and/or large-amplitude photometric variability in the optical; (SE04): very low-mass stars and high-mass brown dwarfs in σ Orionis with high-level variability and/or strong $H\alpha$ and Ca-triplet emission; (He05): bright stars in the Ori OB 1b (super-)association using an HK criterion; (Ke05): low-mass stars and high-mass brown dwarfs in σ Orionis with broadened $H\alpha$ emission; (Ca05): K and M stars in σ Orionis with broadened $H\alpha$ emission; (Ol06): low-mass stars and high-mass brown dwarfs in σ Orionis with $K_s L'$ excess; (Ca06b): $1.0\text{--}0.035 M_\odot$ stars and high-mass brown dwarfs in σ Orionis with excess at $8.0\mu\text{m}$; (Ca07a): brown dwarfs in σ Orionis with excess at $5.8\text{--}8.0\mu\text{m}$; (Ca07b): isolated planetary-mass objects in σ Orionis with excess at $5.8\mu\text{m}$.

$V_T - K_s > 2.0$ mag in Table 10. Four bright stars out of the eight moderate and strong X-ray emitters seem to harbour surrounding discs, which supports the scenario of T Tauri and Ae/Be stars.

The Figure 7 shows the variation between the years 1991 and 1994 of the background subtracted count rate in the energy interval $0.24\text{--}2.0\text{ keV}$ of the flux of the young stars found in WGACAT. Several targets display flares and variability of the X-ray emission in time scales from days to years. For example, the flux from σ Ori AB (top light-curve in Figure 7) raised from 0.257 ± 0.005 to $0.452 \pm 0.011\text{ ct s}^{-1}$ between 1993 Feb. 26 and 28, while the flux from V1147 Ori decreased in a factor 5 with respect to an hypothetical quiescent level at $\sim 0.009\text{ ct s}^{-1}$ in observations spaced out about one year. Both σ Ori AB and V1147 Ori do not apparently possess discs, so the X-ray variability is probably due to stellar winds and chromo- or magnetospheric effects, respectively.

3.3. Different stellar populations surrounding σ Ori

There are five Hipparcos stars in the investigated area, which HIP identifications and parallax determinations are provided in Table 12. The data may suggest that the stars are located at different heliocentric distances. However, the Hipparcos parallax measurements at more than 100 pc are quite not reliable. Furthermore, in at least three cases the measurements are af-




Fig. 7. X-ray light curves of the brightest σ Orionis stars in the ROSAT WGACAT catalogue.

ected by multiplicity. It is expected that the *GAIA* mission will be able to disentangle distinct populations in the Ori OB1 Association located at different heliocentric distances.

Meanwhile, it is possible to disentangle thanks to the knowledge of the spatial velocities of the young stars. Previous determinations of the radial velocity of σ Ori AB and the mean radial velocity of the σ Orionis cluster ranged in a relatively narrow interval between +25 and +35 km s^{-1} (Frost et al. 1926; Heard 1949; Wilson 1953; Evans 1967; Bolton 1974; Cruz-González et al. 1974; Conti et al. 1977; Morrell & Levato 1991; Zapatero Osorio et al. 2002; Muzerolle et al. 2003; Burningham et al. 2005; Kenyon et al. 2005; Caballero 2005). The two recent studies by Jeffries et al. (2006) and Sacco et al. (2006) are in agreement that the heliocentric radial velocity (V_r) is about $+31.0 \pm 0.5 \text{ km s}^{-1}$ with a dispersion less than 8 km s^{-1} (group 2 of Jeffries et al. 2006). I have compiled all the published values of V_r of the investigated stars. They are provided in the last column of Table A.2. Among the nine stars with measured V_r , there is a non-member star (HD 294269, already discussed in Sections 2.2.3 and 2.3.2). The remaining stars have been classified in this work as members and possible members of the Ori OB 1b Association. σ Ori AB, D and E obviously display V_r similar to the mean value of Jeffries et al. (2006). The same occurs for the classical T Tauri star RX J0539.6–0242 AB, at ~ 15 arcmin to the centre of the cluster. However, the measurements at two different epochs of the radial velocity of the Ae/Be star HD 37699 differ from the mean radial velocity ($+14.8 \pm 2.2 \text{ km s}^{-1}$, Neubauer 1943; $+15 \text{ km s}^{-1}$, Duflot et al. 1995). The radial velocity of HD 294298 is also quite discordant, $V_r = +11.5 \text{ km s}^{-1}$. Both HD 37699 and HD 294298 are closely located (in the northeast outer part of the studied area, in the vicinity of the Horsehead and Alnitak), have similar proper motions (identical within the error bars) and possess discs. HD 294297 ($V_r = +25 \text{ km s}^{-1}$) and HD 294268 ($V_r = +20 \text{ km s}^{-1}$) apparently display different V_r from the mean value of σ Orionis as well.

The eastern star in this sample, also close to the Horsehead Nebula, is HD 294297 (with discordant radial velocity and lithium in absorption; Section 2.5.1). Its proper motion is $\geq 30 \text{ mas a}^{-1}$ larger than that of σ Ori AB and similar to those of probable foreground stars with large tangential velocities (see also Zapatero Osorio et al. 2006). The expected positions in the




Fig. 8. Current (filled stars) and future (boxes) positions of the young stars and candidates. Sizes of the boxes are proportional to the uncertainties.

Table 12. Young stars in the Hipparcos catalogue.

Name	HIP	$\pi \pm \delta\pi$ [mas]	Sp. type
σ Ori A+B+IRS1	26549	2.84 ± 0.91	O9.5V+...
σ Ori D	26551	2.84 ± 0.91	B2V
HD 37699	26694	0.63 ± 1.01	B5Vn
HD 37525 AB	26579	2.55 ± 0.99	B5Vp
HD 37333	26438	2.52 ± 1.08	A1Va

future of HD 294297 and of the other 29 young stars are shown in Figure 8. While most of the stars keep their positions in the sky or float around, in the same period of time HD 294297 would have crossed the Horsehead Nebula and be located out of the 30 arcmin-radius area.

From the existence of such young stars with peculiar spatial velocities, it can be derived that: (i) the astrometric criterion used in Section 2.3.2 to discard stars as possible cluster members could be too conservative (but necessary to minimize the number of contaminant foreground stars in the list of candidate young stars); and (ii) there are young stars in the line of sight of the σ Orionis cluster probably originated in a different star-forming region in the Ori OB 1b Association. The second item supports the hypothesis of the existence of at least two spatially superimposed components around σ Ori exposed by Jeffries et al. (2006). They provided an excellent review of some possible consequences of a population with mixed age and distance. However, given that their spectroscopic studied covered a large area, they investigated young stars that are out of the 30 arcmin-radius and closer to the extremely bright O9.5Ib-type Alnitak star than to σ Ori AB itself. Three out of the proper-motion outlier young stars in this Section are located also in the direction of Alnitak, very close to the Horsehead Nebula. What we see may be, in reality, an overlapping of sub-associations linked to different, but relatively close, star-forming regions. This was firstly noted by Béjar et al. (2004a), who proposed that an overdensity of photometric cluster members to the north-east of σ Orionis could be associated to an hypothetical cluster surround-

ing Alnitak. They also studied the radial distribution of low-mass stars and brown dwarfs in a region around σ Ori, and found that the characteristic radius of the surface density, $\rho(r)$, was $r_0 = 8.8 \pm 0.6$ arcmin (where $\rho(r) = \rho_0 e^{-r/r_0}$ and r is the separation to the cluster centre). Since the spatial distribution of stars (split in different mass intervals) and substellar objects in the cluster are identical (Caballero 2006), the spatial density of stellar members at ~ 26 arcmin ($r = 3r_0$) is about 5 % of the central density. Similar results were found by Sherry et al. (2004). Therefore, the contamination by young stars that do not belong to the σ Orionis cluster in wide searches like this may be significative, especially at large separations to the centre. In particular, the four stars with abnormal spatial velocities are located at 21–28 arcmin to σ Ori AB, which supports this hypothesis. The only way to make much headway against these problems is to obtain precise V_r over a wider area, quantify the number of young contaminants and determine if they belong to any of the sub-groups presented by Jeffries et al. (2006) or to new ones. Multi-epoch observations are mandatory, since bona-fide σ Orionis member spectroscopic binaries could appear as V_r outliers and vice versa. Besides, the binary frequency in the higher mass domain of the cluster seems to be as large as ~ 50 % (Caballero 2005).

3.4. Total mass and mass function from 1.1 to $24 M_\odot$

The masses derived for the stars of this sample are about 20 % less than previous determinations in the literature (e.g. Hunger et al. 1989 and Hernández et al. 2005). The list of the twelve stars of σ Orionis more massive than $3 M_\odot$ basically matches that prepared by Sherry et al. (2004). They derived $\sim 100 M_\odot$ as the mass exclusively contained in them. However, in the present work, the stars with $M > 3 M_\odot$ account for only $\sim 64 M_\odot$, which may lead to a cluster total mass significantly lower than previously estimated ($225 \pm 30 M_\odot$; Sherry et al. 2004). Simply by scaling the correction factor down to lower masses, the cluster total mass could be at the level of only $\sim 150 M_\odot$. The total mass could be even lower accounting for probable young contaminants.

The total mass of the stars in this sample (probably all the σ Orionis stars more massive than $\sim 1.2 M_\odot$ plus some young contaminant) is $100^{+20}_{-19} M_\odot$. Out of it, 46 % is contained in the σ Ori A+B+C+D+E system, and up to 29 % only in the two most massive stars in the cluster, σ Ori A and B. This fact indicates that the cluster presents a well-defined cluster centre, in contrast to other extended star-forming regions, as Upper Scorpius (de Zeeuw et al. 1999; Preibisch & Zinnecker 1999). Discarding the three stars discussed in the Section 3.3, the cluster barycentre is located at ~ 1 arcmin to the west of σ Ori AB. The mean tangential velocity of the cluster centre of mass is $(\mu_\alpha \cos \delta, \mu_\delta) = (+2.2 \pm 1.2, -0.5 \pm 1.0) \text{ mas a}^{-1}$, which is also quite similar to the tangential velocity of σ Ori AB (the result rejecting the three stars is identical within the error bars)⁴.

Last, but not less important, I have also computed the mass spectrum in σ Orionis in the mass interval 1.1– $24 M_\odot$ from the masses shown in Table 9. The Figure 9 displays the cumulative number of stars per mass interval (ΔN), the mass spectrum ($\Delta N / \Delta M = \alpha M^\gamma$; $\alpha \equiv -\gamma$) and the mass function ($\Delta \log N / \Delta \log M = \alpha M^\Gamma$; $\alpha \equiv 1 - \Gamma$) in the cluster. The value of the α index from the fit of the mass spectrum for 3 Ma and 360 pc gave $\alpha = +2.0$ (the fit rejecting the four stars in the Section 3.3




Fig. 9. The cumulative number of stars per mass interval (top), the mass spectrum (middle) and the mass function (bottom) in σ Orionis in the mass interval 1.1– $24 M_\odot$. A distance of 360 pc and an age of 3 Ma have been used. In the middle window, the dashed (red) line marks the fit of the mass spectrum ($\alpha = +2.0$).

offers the same result). Considering the very conservative uncertainties in the age and heliocentric distance, and repeating the fit for the less probable cases (1 Ma and 300 pc, 10 Ma and 430 pc), then the value of the index in the mass spectrum is $\alpha = +2.0^{+0.2}_{-0.1}$. This value is slightly smaller than, but quite consistent with, the α value for massive and solar-like stars ($\alpha \approx +2.3$; Salpeter 1955). It may be due to the incompleteness of the search below $1.3 M_\odot$ ($V_T \gtrsim 11.5$ mag). However, the measured α nicely extrapolates the Sherry et al. (2004) studies in σ Orionis. They found that the cluster mass function in the mass interval 0.2– $1.0 M_\odot$ is also consistent with that of the field stars in the same mass interval by Kroupa (2001, 2002). Therefore, the index of the mass spectrum in σ Orionis keeps roughly constant at $\alpha \sim 2$ from 24 to $0.2 M_\odot$, where there is a turning point (first mentioned by Caballero 2006). The α index below $0.2 M_\odot$ is established to be in the range $+0.6$ – 0.8 (Béjar et al. 2001; González-García et al. 2006). Tej et al. (2002), who measured the mass spectrum in an interval covering both sides of the turning point (0.5 – $0.08 M_\odot$), obtained an intermediate value of $\alpha = +1.2 \pm 0.1$. Then, the σ Orionis cluster has become an excellent laboratory to study the mass function from the most massive stars down to planetary-mass objects.

4. Summary

This work is devoted to identify and characterize the brightest stars of the σ Orionis cluster. The results presented here will complement searches of low-mass stars, brown dwarfs and planetary-mass objects in the cluster.

From a correlation between the Tycho-2 and 2MASS catalogues, and using additional information from X-ray and mid-infrared space missions and from the literature, I have classified the brightest stars in a circular area of radius 30 arcmin centred in σ Ori AB in: (i) members of the Ori OB 1b Association with youth features (26); (ii) candidate young stars (4); and (iii) probable foreground stars with red $V_T - K_s$ colours and/or large tangential velocities (16).

⁴ The standard deviation with respect to the mean proper motion (not considering HD 37699, HD 294298 and HD 294297) is 5.1 mas a^{-1} , which transforms into a tangential velocity dispersion of 8.4 km s^{-1} at the heliocentric distance of 360 pc.

The comparison of the sequence of the young stars with the lowest photometric errors and evolutionary tracks in the V vs. $B - V$ diagram suggests that σ Orionis could be closer and older than previously considered.

I have derived the masses from theoretical M_V -mass relations for the young stars and candidate young stars and computed for the first time the mass spectrum of the cluster between 1.1 and 24 $M_\odot$. The α index from the fit of the mass spectrum is $+2.0^{+0.2}_{-0.1}$, which is quite similar to the Salpeter's index. The cluster total mass could be as low as $\sim 150 M_\odot$.

From the flux excess in the mid and/or near infrared, the abnormal position in the V vs. $B - V$ diagram and the strong or moderate X-ray emission, I have derived the existence of discs surrounding seven stars, four out of which are new. The frequency of discs in the magnitude interval $7.5 \text{ mag} \lesssim V \lesssim 11.0 \text{ mag}$ is $32 \pm 14\%$.

Finally, the existence of some stars with abnormal tangential and radial velocities supports the hypothesis of overlapping of stellar populations of different ages and kinematics in the direction of the σ Orionis cluster.

Appendix A: Stellar data

Table A.1 provides the photometric data for the 41 stars in the Tycho-2/2MASS correlation (above the horizontal line) and the five missing stars in the Tycho-2/2MASS correlation (below the horizontal line), ordered by V_T -band magnitude. It is given the name of each star (if not available, then the Tycho identification is provided), the B_T - and V_T -band magnitudes from the Tycho-2 catalogue and the J -, H - and K_s -band magnitudes from the 2MASS catalogue.

Table A.2 provides the astrometric data and the spectral types in the same manner as in Table A.1. It is given the name of each star, right ascension and declination from the Tycho-2 catalogue (FK5, Equinox=J2000.0, Epoch=J2000.000, proper motions taken into account), proper motions in right ascension and in declination from the Tycho-2 catalogue and spectral types from the literature (references in the text).

Acknowledgements. I thank C. Bertout, D. J. Lennon, T. J. Mahoney, V. J. Sánchez Béjar, E. Solano, M. R. Zapatero Osorio and, especially, the referee, R. D. Jeffries, for providing me helpful comments at different stages of the writing. Partial financial support was provided by the Spanish Ministerio de Ciencia y Tecnología project AYA2004-00253 of the Plan Nacional de Astronomía y Astrofísica. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

References

Adams N. R., Wolk S. J., Walter F. M. & Sherry W. H. 2004, American Astronomical Society Meeting 205, #105.08
 Alcalá J. M., Terranegra L., Wichmann R., Chavarría-K. C., Krautter J., Schmitt J. H. M. M., Moreno-Corral M. A., de Lara E.; & Wagner R. M. 1996, A&AS, 119, 7
 Alcalá J. M., Covino E., Torres G., Sterzik M. F., Pfeiffer M. J. & Neuhauser R. 2000, A&A, 353, 186
 Appenzeller I. & Mundt R. 1989, A&ARv, 1, 291
 Barrado y Navascués D., Zapatero Osorio M. R., Béjar V. J. S., Rebolo R., Martín E. L., Mundt R. & Bailer-Jones C. A. L. 2001, A&A, 377, L9
 Barrado y Navascués D., Béjar V. J. S., Mundt R., Martín E. L., Rebolo R., Zapatero Osorio M. R. & Bailer-Jones C. A. L. 2003, A&A, 404, 171
 Béjar V. J. S., Zapatero Osorio M. R. & Rebolo R. 1999, ApJ, 521, 671
 Béjar V. J. S., Martín E. L., Zapatero Osorio M. R., Rebolo R., Barrado y Navascués D., Bailer-Jones C. A. L., Mundt R., Baraffe I., Chabrier C. & Allard F. 2001, ApJ, 556, 830

Béjar V. J. S., Caballero J. A., Rebolo R., Zapatero Osorio M. R. & Barrado y Navascués D. 2004a, Ap&SS, 292, 339
 Béjar V. J. S., Zapatero Osorio M. R. & Rebolo R. 2004b, AN, 325, 705
 Bertout C. 1989, ARA&A, 27, 351
 Blaauw A. 1964, ARA&A, 2, 213
 Blaauw A. 1991, The Physics of Star Formation and Early Stellar Evolution, NATO Advanced Science Institutes (ASI) Series C, Vol. 342, held in Agia Pelagia, Crete, Greece, May 27th - June 8th, Dordrecht: Kluwer, 1991, edited by Charles J. Lada and Nikolaos D. Kylafis., p.125
 Bolton C. T. 1974, ApJ, 192, L7
 Bonnarel F., Fernique P., Bienaymé O., Egret D., Genova F., Louys M., Ochsenein F., Wenger M. & Bartlett J. G. 2000, A&AS, 143, 33
 Briceño C., Calvet N., Hernández J., Vivas A. K., Hartmann L., Downes J. J. & Berlind P. 2005, AJ, 129, 907
 Brown A. G. A., de Geus E. J. & de Zeeuw P. T. 1994, A&A, 289, 101
 Brown A. G. A., Walter F. M. & Blaauw A. 1999, in ASP Conf. Ser., The Orion Complex Revisited, ed. M. J. McCaughrean & A. Burkert, in press
 ten Brummelaar T., Mason B. D., McAlister H. A., Roberts L. C. Jr., Turner N. H., Hartkopf W. I. & Bagnuolo W. G. Jr. 2000, AJ, 119, 2403
 Burningham B., Naylor T., Littlefair S. P. & Jeffries R. D. 2005, MNRAS, 356, 1583
 Caballero J. A., Béjar V. J. S., Rebolo R. & Zapatero Osorio M. R. 2004, A&A, 424, 857
 Caballero J. A. 2005, Astron. Nachr., 326, No. 10, 1007
 Caballero J. A. 2006, PhD thesis, Universidad de La Laguna
 Caballero J. A., Martín E. L., Dobbie P. D. & Barrado y Navascués 2006a, A&A, 460, 635
 Caballero J. A., Barrado y Navascués, Béjar V. J. S., Bouy H., Martín E. L., Rebolo R., Zapatero Osorio M. R. & Bihain G. 2006b, in The 14th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, 6–10 Nov 2006, Pasadena, CA, poster #39
 Caballero J. A. et al. 2007a, submitted to A&A
 Caballero J. A. et al. 2007b, to be submitted to A&A
 Catalano F. A. & Renson P. 1998, A&AS, 127, 421
 Conti P. S., Leep E. M. & Lorre J. J. 1977, ApJ, 214, 759
 Cruz-González C., Recillas-Cruz E., Costero R., Peimbert M. & Torres-Peimbert S. 1974, RMxAA, 1, 211
 Cunha K., Smith V. V. & Lambert D. L. 1995, ApJ, 452, 634
 Cutri R. M., Skrutskie M. F., van Dyk S., Beichman C. A., Carpenter J. M., Chester T., Cambresy L., Evans T., Fowler J., Gizis J. et al. 2003, VizieR On-line Data Catalog: II/246. Originally published in: University of Massachusetts and Infrared Processing and Analysis Center (IPAC/California Institute of Technology)
 Downes R. A. & Keyes C. D.E. 1988, AJ, 96, 777
 Duflot M., Figon P. & Meyssonnier N. 1995, A&AS, 114, 269
 European Space Agency, The Hipparcos and Tycho catalogues, ESA SP-1200, 1997
 Evans D. S. 1967, Determination of Radial Velocities and their Applications, Proceedings from IAU Symposium no. 30 held at the University of Toronto 20–24 June, 1966. Edited by A. H. Batten and J. F. Heard, Academic Press, London, p.57
 Fernique P., Bonnarel F., Louys M., Oberto A., Boch T., Allen M., Bienaymé O. & Egret D. 2004, Toward an International Virtual Observatory: Proceedings of the ESO/ESA/NASA/NSF Conference Held at Garching, Germany, 10–14 June 2002, ESO Astrophysics Symposia. ISBN 3-540-21001-6. Edited by P. J. Quinn and K. M. Górski. Springer-Verlag Berlin/Heidelberg, 2004, p. 271
 Franciosi E., Pallavicini R. & Sanz-Forcada J. 2006, A&A, 446, 501
 Frost E. B., Barrett S. B. & Struve O. 1926, ApJ, 64, 1
 García-Lario P., Manchado A., Suso S. R., Pottasch S. R. & Olling R. 1990, A&AS, 82, 497
 García-Lario P., Manchado A., Pych W. & Pottasch S. R. 1997, A&AS, 126, 479
 Garrison R. F. PASP, 1967, 79, 433
 Genzel R. & Stutzki J. 1989, A&A, 27, 41
 de Geus E. J., Lub J. & van de Grift E. 1990, A&AS, 85, 915
 Gieseke F. 1983, A&A, 118, 102
 González-García B. M., Zapatero Osorio M. R., Béjar V. J. S., Bihain G., Barrado y Navascués D., Caballero J. A. & Morales-Calderón M. 2006, A&A, 460, 799
 Gray R. O. & Corbally C. J. 1993, AJ, 106, 632
 Greenstein J. L. & Wallerstein G. 1958, ApJ, 127, 237
 Gregorio-Hetem J. & Hetem A. Jr. 2000, Planetary Systems in the Universe, International Astronomical Union. Symposium no. 202. Manchester, England, August 2000, meeting abstract 79
 Goudis C., 1982, The Orion complex: A case study of interstellar matter, Dordrecht, Netherlands, D. Reidel Publishing Co. (Astrophysics and Space Science Library. Volume 90), 1982
 Guetter H. H. 1981, AJ, 86, 1057

- Haro G. & Moreno A. 1953, BOTT, 1g, 11
- Hardie R. H., Heiser A. M. & Tolbert C. R. 1964, ApJ, 140, 1472
- Heard J. F. 1949, ApJ, 109, 185
- Hernández J., Calvet N., Hartmann L., Briceño C., Sicilia-Aguilar A. & Berlind P. 2005, AJ, 129, 856
- Hillenbrand L. A., Strom S. E., Vrba F. J. & Keene J. 1992, ApJ, 397, 613
- Høg E., Kuzmin A., Bastian U., Fabricius C., Kuimov K., Lindegren L., Makarov V. V. & Roeser S. 1998, A&A, 335, L65
- Høg E., Fabricius C., Makarov V. V., Urban S., Corbin T., Wycoff G., Bastian U., Schwekendiek P. & Wicenec A. 2000, A&A, 355, L27
- Hunger K., Heber U. & Groote D. 1989, A&A, 224, 57
- Jayawardhana R., Ardila D. R. & Stelzer B. 2003, Brown Dwarfs, Proceedings of IAU Symposium #211, held 20–24 May 2002 at University of Hawai'i, Honolulu, Hawai'i. Edited by E. L. Martín. San Francisco: Astronomical Society of the Pacific, 2003, p. 139
- Jeffries R. D., Maxted P. F. L., Oliveira J. M. & Naylor T. 2006, MNRAS, 371, L6
- Kenyon M. J., Jeffries R. D., Naylor T., Oliveira J. M. & Maxted P. F. L. 2005, MNRAS, 356, 89
- Kharchenko N. V., Piskunov A. E., Röser S., Schilbach E. & Scholz R.-D. 2005, A&A, 440, 403
- Kogure T., Yoshida S., Wiramihardja S. D., Nakano M., Iwata T. & Ogura K. 1989, PASJ, 41, 1195
- Königl A. 1991, ApJ, 370, L39
- Kraemer K. E., Shipman R. F., Price S. D., Mizuno D. R., Kuchar T. & Carey S. J. 2003, AJ, 126, 1423
- Kroupa P. 2001, MNRAS, 322, 231
- Kroupa P. 2002, Science, 295, 82
- Lee T. A. 1968, ApJ, 152, 913
- Lejeune T. & Schaerer D. 2001, A&A, 366, 538
- van Loon J. Th. & Oliveira J. M. 2003, A&A, 405, L33
- Lyngå G. 1981, The Catalogue of Open Star Clusters, ADCBu, 1, 90
- Martín E. L., Zapatero Osorio M. R., Barrado y Navascués D., Béjar V. J. S. & Rebolo R. 2001, ApJ, 558, L117
- Meynet G., Maeder A., Schaller G. & Charbonnel C. 1994, A&AS, 103, 97
- Monet D. G., Levine S. E., Casian B., Ables H. D., Bird A. R., Dahn C. C., Guetter H. H., Harris H. C., Henden A. A., Leggett S. K. et al. 2003, AJ, 125, 984
- Morrell N. & Levato H. 1991, ApJS, 75, 965
- Muzerolle J., Hillenbrand L., Calvet N., Briceño C. & Hartmann L. 2003, ApJ, 592, 266
- Nakano M., Wiramihardja S. D. & Kogure T. 1995, PASJ, 47, 889
- Nesterov V. V., Kuzmin A. V., Ashimbaeva N. T., Volchkov A. A., Röser S. & Bastian U. 1995, A&AS, 110, 367
- Neubauer F. J. 1943, ApJ, 97, 300
- Nissen P. E. 1976, A&A, 50, 343
- Oganesyan R. Kh., Epemyan R. A. & Karapetyan A. A. 1995, Ap, 38, 189
- Oliveira J. M., Jeffries R. D., Kenyon M. J., Thompson S. A. & Naylor T. 2002, A&A, 382, L22
- Oliveira J. M., Jeffries J. D. & van Loon J. Th. 2004, MNRAS, 347, 1327
- Oliveira J. M. & van Loon J. Th. 2004, A&A, 418, 663
- Oliveira J. M., Jeffries R. D., van Loon J. Th. & Rushton M. T. 2006, MNRAS, 369, 272
- Perryman M. A. C., Lindegren L., Kovalevsky J., Hoeg E., Bastian U., Bernacca P. L., Crézé M., Donati F., Grenon M., van Leeuwen F., van der Marel H., Mignard F., Murray C. A., Le Poole R. S., Schrijver H., Turon C., Arenou F., Froeschlé M. & Petersen C. S. 1997, A&A, 323, L49
- Preibisch T. & Zinnecker H. 1999, AJ, 117, 2318
- Reipurth Bo, Bally J., Fesen R. A. & Devine, D. 1998, Nature, 396, 343
- ROSAT Scientific Team 2000a, ROSAT News, 71
- ROSAT Scientific Team 2000b, ROSAT News, 72
- Sacco G. G., Randich S., Franciosini E., Pallavicini R. & Palla F. 2006, A&A preprint doi <http://dx.doi.org/10.1051/0004-6361:20066434>
- Salpeter E. E. 1955, ApJ, 121, 161
- Sanz-Forcada J., Franciosini E. & Pallavicini R. 2004, A&A, 421, 715
- Schaller G., Schaerer D., Meynet G. & Maeder A. 1992, A&AS, 96, 269
- Schild R. E. & Chaffee F. 1971, ApJ, 169, 529
- Scholz A. & Eislöffel J. 2004, A&A, 419, 249
- Sherry W. H., Walter F. M. & Wolk S. J. 2004, AJ, 128, 2316
- Stephenson C. B. 1986, AJ, 92, 139
- Tej A., Sahu K. C., Chandrasekhar T. & Ashok N. M. 2002, ApJ, 578, 523
- Torres C. A. O., Quast G., de La Reza R., Gregorio-Hetem J. & Lepine J. R. D. 1995, AJ, 109, 2146
- Voges W., Aschenbach B., Boller Th., Bräuninger H., Briel U., Burkert W., Dennerl K., Englhauser J., Gruber R., Haberl F. et al. 1999, A&A, 349, 389
- Walter F. M., Wolk S. J., Freyberg M. & Schmitt J. H. M. M. 1997, MmSAI, 68, 108
- Walter F. M., Wolk S. J. & Sherry W. 1998, ASP Conf. Ser., 154, CD-1793
- Warren W. H. Jr. & Hesser J. E. 1977a, ApJS, 34, 115
- Warren W. H. Jr. & Hesser J. E. 1977b, ApJS, 34, 207
- Warren W. H. Jr. & Hesser J. E. 1978, ApJS, 36, 497
- Weaver Wm. B. & Babcock A. 2004, PASP, 116, 1035
- White N. E., Giommi P. & Angelini L. 2001, VizieR On-line Data Catalog: IX/31. Originally published in: Laboratory for High Energy Astrophysics (LHEA/NASA), Greenbelt (2000)
- Wilson R. E. 1953, QB, 901, W495, General catalogue of stellar radial velocities. Washington, [Carnegie Institution of Washington] 1953
- Wiramihardja S. D., Kogure T., Yoshida S., Ogura K. & Nakano M. 1989, PASJ, 41, 155
- Wiramihardja S. D., Kogure T., Yoshida S., Nakano M., Ogura K. & Iwata T. 1991, PASJ, 43, 27
- Wolk S. J. 1996, Ph.D. thesis, State Univ. New York at Stony Brook
- Wolk S. J. & Walter F. M. 2000, Very Low-mass Stars and Brown Dwarfs, ed. R. Rebolo & M. R. Zapatero Osorio, 38
- Zacharias N., Monet D. G., Levine S. E., Urban S. E., Gaume R. & Wycoff G. L. 2004, American Astronomical Society Meeting 205, #48.15; Bulletin of the American Astronomical Society, Vol. 36, p. 1418
- Zapatero Osorio M. R., Béjar V. J. S., Martín E. L., Rebolo R., Barrado y Navascués D., Bailer-Jones C. A. L. & Mundt R. 2000, Science, 290, 103
- Zapatero Osorio M. R., Béjar V. J. S., Pavlenko Ya., Rebolo R., Allende Prieto C., Martín E. L. & García López R. J. 2002, A&A, 384, 937
- Zapatero Osorio M. R., Béjar V. J. S., Bihain G. et al. 2006, A&A, submitted
- de Zeeuw P. T., Hoogerwerf R., de Bruijne J. H. J., Brown A. G. A. & Blaauw A. 1999, AJ, 117, 354

Table A.1. Optical and near-infrared photometry of all the stars.

Name	$B_T \pm \delta B_T$ [mag]	$V_T \pm \delta V_T$ [mag]	$J \pm \delta J$ [mag]	$H \pm \delta H$ [mag]	$K_s \pm \delta K_s$ [mag]
σ Ori A+B+IRS1	3.521 0.014	3.763 0.009	4.752 0.260	4.641 0.252	4.490 0.016
σ Ori E	6.236 0.014	6.344 0.010	6.974 0.026	6.954 0.031	6.952 0.029
HD 37699	7.457 0.015	7.590 0.011	7.841 0.029	7.904 0.042	7.880 0.031
HD 294271	7.757 0.016	7.856 0.011	8.100 0.021	8.180 0.055	8.202 0.038
HD 37525 AB	7.952 0.016	8.058 0.012	8.131 0.030	8.105 0.042	8.093 0.020
HD 294272 A	8.402 0.017	8.389 0.015	8.346 0.026	8.380 0.046	8.374 0.026
HD 294272 B ^a	8.544 0.018	8.554 0.015	8.779 0.021	8.819 0.065	8.790 0.025
HD 37333	8.616 0.017	8.508 0.013	8.413 0.023	8.481 0.065	8.407 0.040
HD 37564	8.699 0.017	8.443 0.014	7.976 0.021	7.881 0.038	7.828 0.029
V1147 Ori	9.065 0.018	8.992 0.016	8.876 0.024	8.894 0.063	8.800 0.019
HD 37686	9.210 0.019	9.186 0.018	9.207 0.035	9.203 0.024	9.155 0.022
HD 37545	9.263 0.018	9.277 0.017	9.261 0.023	9.303 0.024	9.298 0.023
HD 294275	9.504 0.020	9.431 0.019	9.256 0.028	9.242 0.023	9.183 0.025
HD 294268	10.966 0.049	10.507 0.043	9.393 0.023	9.061 0.024	8.873 0.023
HD 294307	11.091 0.054	10.552 0.046	9.439 0.026	9.177 0.023	9.127 0.019
RX J0539.6–0242 AB ^b	11.103 0.055	10.195 0.036	8.462 0.027	8.055 0.040	7.944 0.038
HD 294279	11.131 0.054	10.685 0.053	9.873 0.028	9.727 0.023	9.683 0.026
EM* StHA 50	11.219 0.059	11.275 0.089	10.666 0.024	10.614 0.024	10.545 0.024
GSC 04771–00621	11.231 0.059	10.920 0.063	9.796 0.027	9.600 0.023	9.503 0.027
HD 294277	11.416 0.063	9.737 0.024	6.773 0.017	6.066 0.034	5.861 0.021
HD 294278	11.449 0.067	9.941 0.029	7.592 0.023	7.034 0.053	6.855 0.033
HD 294270	11.507 0.067	10.960 0.062	9.780 0.024	9.511 0.025	9.472 0.025
HD 294301	11.541 0.079	11.139 0.084	10.210 0.028	10.059 0.024	9.992 0.024
HD 294269	11.597 0.076	10.795 0.011	9.178 0.035	8.721 0.033	8.634 0.027
GSC 04771–00962	11.608 0.073	11.101 0.075	10.202 0.026	9.960 0.023	9.905 0.022
HD 294298	11.646 0.086	10.975 0.073	9.339 0.029	8.902 0.061	8.802 0.025
HD 294274	11.662 0.071	10.758 0.056	9.428 0.027	9.069 0.023	8.985 0.025
[W96] 4771–0950 ^c	11.807 0.086	11.429 0.099	10.088 0.027	9.829 0.024	9.750 0.022
HD 294280	11.893 0.091	9.865 0.027	6.429 0.019	5.580 0.031	5.340 0.024
TYC 4771 720 1	11.944 0.105	11.787 0.152	10.269 0.028	9.885 0.023	9.813 0.021
2E 0535.4–0241	12.021 0.117	11.015 0.069	9.255 0.020	8.720 0.051	8.613 0.025
TYC 4770 1261 1	12.232 0.160	11.498 0.106	10.721 0.048	10.458 0.046	10.299 0.042
TYC 4771 661 1	12.386 0.201	11.899 0.178	10.617 0.026	10.298 0.026	10.225 0.024
TYC 4770 1432 1	12.799 0.255	12.337 0.211	11.077 0.023	10.844 0.025	10.731 0.023
TYC 4770 1129 1	12.866 0.240	11.827 0.146	11.232 0.021	11.034 0.024	10.979 0.023
TYC 4770 1018 1	13.014 0.280	11.224 0.077	8.620 0.021	7.881 0.027	7.730 0.017
SO120532 ^d	13.094 0.317	12.031 0.172	11.125 0.026	10.962 0.026	10.835 0.024
TYC 4771 1012 1	13.009 0.302	11.263 0.090	8.998 0.019	8.365 0.061	8.205 0.026
TYC 4771 934 1	13.325 0.387	12.870 0.306	8.948 0.032	8.374 0.057	8.207 0.026
TYC 4771 1468 1	13.433 0.343	11.136 0.074	8.616 0.026	7.893 0.053	7.727 0.023
TYC 4770 924 1	13.367 0.358	11.938 0.163	10.725 0.022	10.317 0.027	10.222 0.025
σ Ori D	6.375 0.012	6.557 0.014	7.116 0.029	7.219 0.027	7.260 0.021
σ Ori C ^e	8.77 0.02	8.79 0.02	9.086 0.032	9.109 0.047	9.129 0.021
HD 294273	10.899 0.106	10.828 0.186	10.176 0.028	10.099 0.026	10.103 0.027
HD 294297	10.905 0.090	10.193 0.073	9.062 0.032	8.872 0.084	8.775 0.024
HD 294276	11.193 0.071	10.418 0.057	9.183 0.021	8.856 0.025	8.814 0.026

^a Also BD–02 1323C.^b Also 2E 0537.1–0243.^c Name from Wolk (1996). Also SO210457.^d Name from Caballero (2006).^e Johnson *B*- and *V*-band photometry from Greenstein & Wallerstein (1958)

Table A.2. Astrometry, spectral type and radial velocity (when available) of all the stars.

Name	TYC	α (ICRS)	δ (ICRS)	$\mu_\alpha \cos \delta$ $\pm \delta \mu_\alpha \cos \delta$ [mas a ⁻¹]	μ_δ $\pm \delta \mu_\delta$ [mas a ⁻¹]	Sp. type	V_r km s ⁻¹
σ Ori A+B+IRS1 ^a	4771 1196 1	05 38 44.765	-02 36 00.25	+4.61±0.88	-0.40±0.53	O9.5V+...	+29.1
σ Ori E	4771 1194 1	05 38 47.208	-02 35 40.52	+2.2±1.1	-1.4±1.2	B2Vp	+29
HD 37699	4771 1045 1	05 40 20.189	-02 26 08.22	+0.3±1.2	+0.9±1.3	B5Vn	+14.8±2.2
HD 294271	4771 1193 1	05 38 36.549	-02 33 12.76	+0.4±1.3	-0.6±1.4	B5V	
HD 37525 AB	4771 1103 1	05 39 01.493	-02 38 56.36	+2.1±1.3	-0.4±1.3	B5Vp	
HD 294272 A	4771 1205 1	05 38 34.799	-02 34 15.78	-3.3±1.6	+0.6±1.6	B9.5III	
HD 294272 B	4771 1204 1	05 38 34.235	-02 34 16.08	+5.4±1.8	-3.4±1.7	B8V	
HD 37333	4771 956 1	05 37 40.476	-02 26 36.83	-3.3±1.0	-3.9±1.3	A1Va	
HD 37564	4771 1073 1	05 39 15.061	-02 31 37.62	+2.2±1.2	+0.5±1.2	A8V:	
V1147 Ori	4771 685 1	05 39 46.196	-02 40 32.06	-2.5±1.2	+4.0±1.3	B9IIIp	
HD 37686	4771 909 1	05 40 13.090	-02 30 53.18	+2.1±1.2	+2.1±1.2	B9.5Vn	
HD 37545	4771 304 1	05 39 09.213	-02 56 34.73	+1.9±1.1	-1.1±1.1	B9V	
HD 294275	4771 39 1	05 37 31.872	-02 45 18.48	-2.7±1.2	-4.4±1.3	A1V	
HD 294268	4771 802 1	05 38 14.121	-02 15 59.78	+4.5±1.5	+0.8±1.5	F5	+20
HD 294307	4771 157 1	05 40 12.462	-02 52 57.59	+0.1±1.4	-16.8±1.4	F8	
RX J0539.6-0242 AB	4771 543 1	05 39 36.541	-02 42 17.21	+2.2±1.5	+2.8±1.6	G5-K0	+32.6±1.8
HD 294279	4771 405 1	05 38 31.382	-02 55 03.15	-0.4±1.4	+6.7±1.4	F3-5	
StHA 50	4771 1087 1	05 38 34.449	-02 28 47.56	+2.8±1.6	+3.2±1.6	Be	
GSC 04771-00621	4771 621 1	05 37 41.793	-02 29 08.21	+7.0±1.6	-11.1±1.6		
HD 294277	4771 463 1	05 37 57.342	-02 53 17.69	+4.8±1.2	-2.8±1.2	K2	
HD 294278	4771 127 1	05 38 38.762	-02 49 01.31	+10.9±1.4	+2.6±1.4	K2	
HD 294270	4770 1336 1	05 37 18.818	-02 31 36.44	+16.8±1.9	-23.4±1.8	G0	
HD 294301	4771 556 1	05 40 21.125	-02 40 25.62	+4.6±1.7	-8.5±1.7	F2V(n)	
HD 294269	4771 693 1	05 37 57.818	-02 26 33.65	+43.5±1.8	-16.1±1.8	G0	+72.8
GSC 04771-00962	4771 962 1	05 37 44.915	-02 29 57.31	+2.8±1.9	+2.7±2.1		
HD 294298	4771 880 1	05 39 59.318	-02 22 54.35	-0.4±1.8	+1.8±1.8	G0:	+11.5
HD 294274	4771 940 1	05 37 45.365	-02 44 12.53	-8.1±3.2	-19.4±3.5	G0	
4771-0950	4771 950 1	05 38 06.496	-02 28 49.37	+6.9±1.8	-7.8±1.8	F7	
HD 294280	4771 385 1	05 38 28.491	-03 03 33.79	-4.0±1.7	+4.9±1.8	K5	
TYC 4771 720 1	4771 720 1	05 37 59.046	-02 41 00.47	-20.1±3.9	-16.6±4.2		
2E 0535.4-0241	4771 921 1	05 37 54.405	-02 39 29.84	-1.8±3.8	+0.9±4.0		
TYC 4770 1261 1	4770 1261 1	05 37 10.470	-02 30 07.19	-4.3±3.8	-8.5±4.1		
TYC 4771 661 1	4771 661 1	05 39 20.446	-02 27 51.44	+16.6±4.3	-20.1±4.7		
TYC 4770 1432 1	4770 1432 1	05 37 09.689	-02 39 59.87	-30.9±4.5	+0.1±5.0		
TYC 4770 1129 1	4770 1129 1	05 37 08.495	-02 22 06.34	-5.8±4.2	+0.4±4.7		
TYC 4770 1018 1	4770 1018 1	05 36 57.142	-02 25 40.00	-4.3±4.2	-2.3±4.6		
SO120532	4771 873 1	05 38 14.438	-02 19 58.64	-3.0±4.3	-5.5±4.7	F7-9	
TYC 4771 1012 1	4771 1012 1	05 39 13.944	-02 10 49.27	-0.9±4.2	-9.7±4.6		
TYC 4771 934 1	4771 934 1	05 39 43.058	-02 28 45.59	-7.5±4.6	-2.7±4.9		
TYC 4771 1468 1	4771 1468 1	05 37 29.879	-02 43 45.83	-7.0±4.7	-6.5±4.9		
TYC 4770 924 1	4770 924 1	05 37 19.240	-02 28 46.22	+1.8±4.7	-43.4±5.1		
σ Ori D ^b	4771 1195 1	05 38 45.602	-02 35 58.84	+4.6±0.9	-0.4±0.5	B2V	+33.1±3.1
σ Ori C ^c	—	05 38 44.12	-02 36 06.3	0±0	0±0	A2V	
HD 294273 ^b	4771 1023 1	05 38 27.524	-02 43 32.60	-0.6±2.8	+4.6±2.5	A3	
HD 294297 ^b	4771 1081 1	05 40 27.531	-02 25 42.90	+22.7±2.6	-20.5±1.7	F6-8	+25
HD 294276 ^b	4770 1489 1	05 37 20.671	-02 49 32.55	+8.8±1.7	-61.7±3.6	G0	

^a Proper motions from Perryman et al. (1997). The spectral types of the components of the triple system are O9.5V, B0.5V and K:.^b Proper motions from Høg et al. (1998).^c Coordinates from the 2MASS catalogue and proper motion from the NOMAD1 catalogue.

List of Objects

'Alnitak' on page 1	'HD 294276' on page 14
'Alnilam' on page 1	'BD-02 1323C' on page 14
'Mintaka' on page 1	'2E 0537.1-0243' on page 14
'Horsehead Nebula' on page 1	'SO210457' on page 14
'Flame Nebula' on page 1	
'NGC 2024' on page 1	
'IC 434' on page 1	
'SE 70' on page 2	
'S Ori 68' on page 2	
'SE 70' on page 2	
' σ Ori A' on page 2	
' σ Ori B' on page 2	
' σ Ori IRS1' on page 2	
'HD 37744' on page 3	
'TX Ori' on page 3	
'V510 Ori' on page 3	
'IRAS 05358-0238' on page 3	
'IRAS 05352-0227' on page 3	
' σ Ori E' on page 14	
'HD 37699' on page 14	
'HD 294271' on page 14	
'HD 37525' on page 14	
'HD 294272 A' on page 14	
'HD 294272 B' on page 14	
'HD 37333' on page 14	
'HD 37564' on page 14	
'V1147 Ori' on page 14	
'HD 37686' on page 14	
'HD 37545' on page 14	
'HD 294275' on page 14	
'HD 294268' on page 14	
'HD 294307' on page 14	
'RX J0539.6-0242' on page 14	
'HD 294279' on page 14	
'EM* StHA 50' on page 14	
'GSC 04771-00621' on page 14	
'HD 294277' on page 14	
'HD 294278' on page 14	
'HD 294270' on page 14	
'HD 294301' on page 14	
'HD 294269' on page 14	
'GSC 04771-00962' on page 14	
'HD 294298' on page 14	
'HD 294274' on page 14	
'[W96] 4771-0950' on page 14	
'HD 294280' on page 14	
'TYC 4771 720 1' on page 14	
'2E 0535.4-0241' on page 14	
'TYC 4770 1261 1' on page 14	
'TYC 4771 661 1' on page 14	
'TYC 4770 1432 1' on page 14	
'TYC 4770 1129 1' on page 14	
'TYC 4770 1018 1' on page 14	
'SO120532' on page 14	
'TYC 4771 1012 1' on page 14	
'TYC 4771 934 1' on page 14	
'TYC 4771 1468 1' on page 14	
'TYC 4770 924 1' on page 14	
' σ Ori D' on page 14	
' σ Ori C' on page 14	
'HD 294273' on page 14	
'HD 294297' on page 14	